

Effect of selected treatments on vase-life and freshness of rose flowers in home decoration

■ JAYSHREE RODGE

Author for correspondence:

JAYSHREE RODGE

Assistant Professor
Department of Family
Resource Management
College of Home Science
Maharashtra Agricultural
University, PARBHANI
(M.S.) INDIA

ABSTRACT

For studying in roses with stem, various concentrations of solutions used were Filtered water-control (T_0), Sucrose – 1% (T_1), Sodium chloride-1% (T_2), Gibberlic acid – 60 ppm (T_3) and Glycerin -5 ml / l. Two sets of all the treatments were used to assess the effect of daily addition of the solutions in the freshness of roses. In one set (V_0) the respective solutions were added daily to initial level whereas the other set (V_1) was kept without addition. Treatment with 1 per cent salt (T_2) extended the vase-life of roses and had significantly better colour, freshness and firmness of stems. Every day addition of respective solutions to make up the absorbed volume extended the vase-life and had better floral appearance. Similarly, total rate of solution uptake was high and the fresh weight was also more when the volume was made up.

KEY WORDS : Roses, Vase-life, Freshness, Home decoration

How to cite this paper: Rodge, Jayshree (2011). Effect of selected treatments on vase-life and freshness of rose flowers in home decoration. *Asian J. Home Sci.*, 6 (2) : 284-286.

Article chronicle: Received: 08.06.2011; **Revised:** 08.10.2011; **Accepted:** 30.11.2011

Flowers, the nature's bounty, are loved, admired and adored. Available in fascinating colours, forms and shapes, they express happiness, cheerfulness, reverence, on all occasions of social, cultural and religious functions.

The life span and freshness of flowers depend upon the factors such as rate of absorption and transpiration, season, cutting time (Devlin, 1968; Larmine, 1980). Prolonging the life span of flowers, after separating from mother plant, is attempted in laboratories and certain treatments have been successfully developed for commercial purpose and various materials have been recommended such as charcoal, sugar, common salt, aspirin, olive oil, camphor (Thamman, 1980; Mullik, 1981; Khanna, 1982). Preserving roses with glycerin helps them to maintain more of their colour, shape and texture than normal dried roses have so (Kitty Belendez, 2011).

Validity of such recommended treatments therefore, needs to be tested to help these assets. The present study has therefore been conducted with the following objectives : to find out the effect of selected household treatments in terms of vase-life and freshness of rose flowers and to find out the effect of daily addition of the solutions to make up the absorbed volume in retaining the freshness of rose flowers.

RESEARCH METHODS

For the experiment, the roses with stem were

selected as these have been found to be used commonly. The flower was standardized for the stem length, size and weight, freshness in appearance.

For studying the roses with stem, sucrose, common salt were the materials selected (a recommended by Larmie, 1980 and Mullik, 1981). While gibberlic acid and glycerine were included as effective laboratory treatments. Standardization of the materials was done in respect to the quality of water, concentration and the initial amount to be used.

The length of roses was kept 7 inches and weight ranged between 10 to 15 g. Concentration of solutions used were: Filtered used –control (T_0), Sucrose-1% (T_1), Sodium chloride-1% (T_2), Gibberlic acid – 60 ppm (T_3) and Glycerine -5 ml/l (T_4).

Two sets of all the treatments was used to assess the effect of daily addition of the solutions in the freshness of roses. In one set (V_0) the respective solutions were added daily to initial level where as the other set (V_1) was kept without addition.

RESEARCH FINDINGS AND DISCUSSION

It is depicted from the results of Table 1 that the mean vase-life of roses treated with 1 per cent salt (T_2) was longer (3.83 days) compared to control (3 days) and all other treatments. The same treatments (T_2) secured highest mean score for six selected parameters compared