

Research
Paper

Effect of sowing dates on growth and yield of French bean (*Phaseolus vulgaris* L.) varieties during *Kharif* season

MAMTA J. PATANGE, N.G. LAD AND SHUBHANGI J. DHAGE

See end of the paper for authors' affiliations

Correspondence to :

MAMTA J. PATANGE
Agricultural Technical School
(M.A.U.), LATUR (M.S.)
INDIA

ABSTRACT

An experiment was conducted during *Kharif* season at Department of Agronomy, Marathwada Agricultural university, Parbhani to study the effect of sowing dates on growth and yield of French bean (*Phaseolus vulgaris* L.). The experiment was conducted in split plot design with 3 replications. Each replication consisted of 16 (sixteen) treatment combinations comprising 04 (four) sowing dates *i.e.* 10 days interval after first sowing on onset of monsoon and four varieties *i.e.* Varun, Waghya, Arka komal and Contender. Sowing date treatment were allotted to the main blocks randomly and varieties were allotted randomly in each sub-block. Results showed, sowing date M_1 (onset of monsoon) and M_2 (10 days after first sowing) were at par with each other and both sowing dates recorded significantly higher seed yield (kg/ha) over M_3 (20 days after first sowing) and M_4 (30 days after first sowing). Sowing date M_3 (20 days after first sowing) was significantly superior over M_4 (30 days after first sowing) which recorded, significantly lowest seed yield (kg/ha). Variety V_1 (Varun) recorded significantly higher grain yield (820kg/ha) over remaining three varieties. However varieties V_4 (contender), V_3 (Arka komal) and V_2 (Waghya) were at par with each other. Variety V_2 (Waghya) recorded significantly lowest seed yield (365kg/ha).

Patange, Mamta J., Lad, N.G. and Dhage, Shubhangi J. (2011). Effect of sowing dates on growth and yield of French bean (*Phaseolus vulgaris* L.) varieties during *Kharif* season, *Adv. Res. J. Crop Improv.*, 2 (2) : 158-160.

KEY WORDS : Sowing dates, Varieties, Yield attributes, Yield, French bean

French bean (*Phaseolus vulgaris* L.) is an important pulse vegetative crop of the world. Brazil, China and United States are the important countries which are producing more than half of the world's supply (Anonymous, 1995). It is probably a native of Southern and central America (Parthasarthy, 1986). But in India it can be grown during *Rabi* season. Time of sowing is a non-monetary input which influences grain yield to a great extent. Optimum time of sowing may vary with the location. Variety may also vary in growth and maturity and thus influence grain yield. Further more, the optimum time of sowing may vary with different varieties of French bean. Therefore, field experiments were conducted to study the performance of varieties of French bean under different dates of sowing.

RESEARCH PROCEDURE

A field experiment was conducted at Department of Agronomy, Marathwada Agricultural university Parbhani, to study the effects of sowing dates on growth and yield of French bean varieties during *Kharif* season of 2005-06. The soil of the experimental field was clayey,

fairly rich in available potassium, low in available nitrogen and medium in phosphorus.

The experiment was laid out in split plot design with three replication. Each replication consisted of 16 treatment combinations comprising 4 sowing dates *i.e.* M_1 - onset of monsoon, M_2 - 10 days after first sowing, M_3 - 20 days after first sowing, M_4 - 30 days after first sowing and four varieties *i.e.* V_1 - Varun, V_2 - Waghya, V_3 - Arka Komal and V_4 - contender. Each replication was divided into four main blocks and each block was divided into sub-blocks. The gross and net plot size was 4.5m x 3.6m and 3.6m x 2.7m, respectively. Sowing date treatments were allotted to the main blocks randomly and varieties were allotted randomly in each sub-block.

RESEARCH ANALYSIS AND REASONING

The results obtained from the present investigation have been discussed below:

Sowing dates:

It was observed from the Table 1 that the sowing date M_1 was significantly superior over sowing date M_3 ,