

Effect of Fe-EDTA on growth, yield and quality of red chilli (*Capsicum annuum* L.)

H.R. SAVITHA¹, B.I. BIDARI¹, G.C. SHASHIDHARA, K.N. KALYANA MURTHY*,
D.S. POORNIMA¹ AND K.S.MOHAN¹

Department of Agronomy, University of Agriculture Science, DHARWAD (KARNATAKA) INDIA

ABSTRACT

An experiment was carried out to study the effect of soil and foliar application of Fe-EDTA on yield and quality of chilli (Cv. Byadgi dabbi) in a calcareous Vertisols at UAS, Dharwad during 2006. Soil application of Fe-EDTA equivalent to FeSO_4 at $20 \text{ kg ha}^{-1} + 0.5\%$ Fe-EDTA foliar spray at 50 DAT was found most effective and recorded significantly higher plant height (83.12 cm), number of branches (17.92), dry matter production ($114.32 \text{ g plant}^{-1}$), dry fruit yield (10.5 q ha^{-1}) and ascorbic acid content ($178.90 \text{ mg } 100\text{g}^{-1}$). But soil application of Fe-EDTA equivalent to FeSO_4 at $20 \text{ kg ha}^{-1} + 0.5\%$ Fe-EDTA foliar spray at 50 and 90 DAT was recorded highest colour value (228.7 ASTA units) and oleoresin content (16.76 %). Where as lowest plant height (70.98 cm), number of branches (12.91), dry matter production ($73.13 \text{ g plant}^{-1}$) dry fruit yield (7.65 q ha^{-1}), ascorbic acid content ($127.6 \text{ mg } 100\text{g}^{-1}$), colour value (163.1 ASTA units) and oleoresin content (13.03 %) were recorded with application of recommended dose of fertilizers.

Key words : Iron-EDTA, Byadgi dabbi, Colour value, Ascorbic acid, Oleoresin, Vertisol

INTRODUCTION

Chilli (*Capsicum annuum* L.) is an important spice cum vegetable crop grown in medium black and deep black calcareous vertisols in Northern Karnataka. In these soils one of the the major problems is, availability of iron is low and it is severe in Karnataka state. In order to correct the deficiency of iron the common practice is to apply Fe either through soil or foliar application. In this respect, if iron is applied in the form of chelate (Fe-EDTA) the efficiency will be high and response to applied iron is quick. It is essential to supply the iron either through soil or foliar spray for obtaining better productivity as well as quality. The price of red chillies is mainly based on the colour value. Wrinkled shining blood red coloured fruits fetch highest price. Among the micronutrients iron appears to play an important role in the synthesis of red colour in chillies (Malawadi *et al.*, 2004). Further, iron content of leaves and stems is closely related to colour value of red chillies (Martinez *et al.*, 1990). Since chilli is being extensively cultivated in calcareous Vertisols in northern Karnataka, it is likely to be deficient in these soils. Information pertaining to the use of Iron-Chelate (Fe-EDTA) as source of iron in improving the quality attributes of chillies grown in calcareous Vertisol is lacking.

MATERIALS AND METHODS

The experiment was carried out during *Kharif* 2006 in a calcareous Vertisols at Main Agricultural Research Station, University of Agricultural Sciences, Dharwad

(Karnataka). The soil pH was 7.40, electrical conductivity 0.42 dSm^{-1} , organic carbon 6.10 g kg^{-1} , free lime content 9.80%, available iron 3.10 mg kg^{-1} , available nitrogen 350.5 kg ha^{-1} , available phosphorus 27.2 kg ha^{-1} , available potassium 410.9 kg ha^{-1} and available sulphur 24.4 ppm. The amorphous form of Fe-EDTA containing 12% iron was used as source of iron. The experiment consisted of twelve treatments replicated thrice and laid out in a Randomized Complete Block Design (RCBD). The Fe-EDTA was applied to soil and as foliar spray. Soil application was done at planting with two levels and foliar application was done at different crop growth stages. The recommended dose of N, P and K were applied @ $100:50:50 \text{ kg ha}^{-1}$ as basal dose through urea, diammonium phosphate and muriate of potash, respectively. The variety of chilli used was Byadgi dabbi. All the cultural practices were followed as per the package of practices of UAS, Dharwad.

RESULTS AND DISCUSSION

The results obtained from the present investigation as well as relevant discussion have been presented under following heads:

Growth and yield:

Soil application of Fe-EDTA resulted in significantly higher growth and yield compared to control (Table 1). Soil application of Fe-EDTA equivalent to FeSO_4 at $20 \text{ kg ha}^{-1} + 0.5\%$ Fe-EDTA as foliar spray at 50 days after transplanting (DAT) recorded as significantly higher plant

* Author for correspondence.

¹Department of Soil Science and Agricultural Chemistry, University of Agricultural Sciences, DHARWAD (KARNATAKA) INDIA