



Research Paper

Article history :

Received : 09.07.2012

Revised : 16.10.2012

Accepted : 16.11.2012

Effect of different grades of rhizomes on growth and yield of turmeric (*Curcuma longa* L.)

■ K. PADMADEVI, L. JEEVA JOTHI¹, V. PONNUSWAMI¹, V. DURGAVATHI¹
AND I. RIJWANA PARVEEN¹

Members of the Research Forum

Associated Authors:

¹Horticultural College and Research Institute, Tamil Nadu Agricultural University, PERIYAKULAM (T.N.) INDIA

Author for correspondence :

K. PADMADEVI

Horticultural College and Research Institute, Tamil Nadu Agricultural University, PERIYAKULAM (T.N.) INDIA
Email : padma_horti@yahoo.co.in

ABSTRACT : Turmeric (*Curcuma longa* L.) plant species produces different sizes of mother rhizomes and finger rhizomes. Rhizomes are used as propagating material in turmeric cultivation. The effects of seed rhizome size on growth and yield of turmeric was evaluated. Mother rhizomes of 10- 15g to 45-50g and finger rhizomes of 5-10 g to 25-30g were tested. The heavier the mother rhizome (45-50 g), better the plant growth. Plants from mother rhizome (25-30g) and finger rhizomes (25-30g) grew similarly well. The seed rhizomes with a greater diameter developed vigorous seedlings. The plants grown from mother rhizome (45-50g) reported the highest plant growth characters like plant height (31.43 cm), number of leaves (7.20), number of tillers (2.81), stem girth (9.14 cm) and leaf area (21012 cm² m⁻²), which were significantly higher than those from lighter finger rhizomes. Finger rhizomes (10-15g) was easily broken at the time planting, and had secondary and tertiary finger rhizomes, which developed thinner plants and resulted in a lower yield. The yield and dry weight were maximum in the plants grown directly from mother rhizome (45-50g) and lower in the plants grown from finger rhizomes (10-15g). This study indicates that the turmeric seed rhizome should be (45-50g) with a larger diameter.

KEY WORDS : Seed rhizome, Mother rhizome, Finger rhizome, Growth, Yield

HOW TO CITE THIS ARTICLE : Padmadevi, K., Jeeva Jothi, L., Ponnuswami, V., Durgavathi, V. and Rijwana Parveen, I. (2012). Effect of different grades of rhizomes on growth and yield of turmeric (*Curcuma longa* L.), *Asian J. Hort.*, 7(2) : 465-467.

Turmeric (*Curcuma longa* L.) is a small perennial herb native to India. It is used as condiment, dye, drug and cosmetic in addition to its use in religious ceremonies. Turmeric belongs to Zingiberaceae family and is cultivated extensively in Asia especially India, China, and other countries. India is a leading producer and exporter of turmeric in the world. In India, Andhra Pradesh, Tamil Nadu, Orissa, Karnataka, West Bengal, Gujarat, Meghalaya, Maharashtra, Assam are the important states which cultivates turmeric, of which, Andhra Pradesh alone occupies 35.0 per cent of area and 47.0 per cent of production. In India, turmeric is cultivated under 1, 80,960 ha with the production of 7, 92,980 MT (NHB database, 2011). The active constituent present in turmeric is curcumin, which comprises 0.3-5.4 per cent (Leung, 1980). Curcuminoids in turmeric have anti-inflammatory, antimutagen, anticancer, antibacterial, anti-oxidant, antifungal, antiparasitic and detoxifying properties (Herrmann and Martine, 1991;

Nakamura *et al.*, 1998; Osawa *et al.*, 1995; Sugiyama *et al.*, 1996 and Uechi *et al.*, 2000).

Normally turmeric is propagated through a small portion of rhizomes known as seed rhizome or seed sets (Dupriez and De leener, 1992; Borget, 1993; Ravindran *et al.*, 2004). The seed rhizome gives the economic yield. The planting material used affects the growth and yield of the crops. Therefore, selecting the right size of planting material (length, weight and number of growing buds per seed) is a very critical factor in the cultivation of turmeric. The use of large seed rhizomes is generally found to increase the final yield of rhizomatous spices such as ginger (Whiley, 1990; Borget, 1993). Large sized seed rhizomes of ginger give significantly higher yield than planting of small pieces (Nybe and Raj, 2004). Hossain *et al.* (2005) found high yield of turmeric from using 30-40 g seed rhizomes compared to 10 and 20 g seed rhizomes. Therefore, the objective of this study was to determine the optimum seed